

10/658904

=> d his

(FILE 'HOME' ENTERED AT 12:20:43 ON 07 APR 2006)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:21:52 ON 07 APR 2006

```
L1      1 S "14171 KINASE?"
L2      1434878 S KINASE?
L3      7629584 S CLON? OR EXPRESS? OR RECOMBINANT
L4      73 S "T-P" MOTIF
L5      49 S L2 AND L4
L6      12 DUP REM L5 (37 DUPLICATES REMOVED)
L7      947418 S (MODULAT? OR ACTIVAT? OR INHIBIT? ) AND L2
L8      42 S "14171"
L9      2 S L7 AND L8
        E LIBERMANN ROSANNA
        E LIBERMANN/AU
        E KAPELLER R/AU
L10     103 S E3
L11     0 S L8 AND L10
```

=>

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal652mxm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	DEC 21	IPC search and display fields enhanced in CA/CAPLUS with the IPC reform
NEWS	4	DEC 23	New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/USPAT2
NEWS	5	JAN 13	IPC 8 searching in IFIPAT, IFIUDB, and IFICDB
NEWS	6	JAN 13	New IPC 8 SEARCH, DISPLAY, and SELECT enhancements added to INPADOC
NEWS	7	JAN 17	Pre-1988 INPI data added to MARPAT
NEWS	8	JAN 17	IPC 8 in the WPI family of databases including WPIFV
NEWS	9	JAN 30	Saved answer limit increased
NEWS	10	JAN 31	Monthly current-awareness alert (SDI) frequency added to TULSA
NEWS	11	FEB 21	STN AnaVist, Version 1.1, lets you share your STN AnaVist visualization results
NEWS	12	FEB 22	Status of current WO (PCT) information on STN
NEWS	13	FEB 22	The IPC thesaurus added to additional patent databases on STN
NEWS	14	FEB 22	Updates in EPFULL; IPC 8 enhancements added
NEWS	15	FEB 27	New STN AnaVist pricing effective March 1, 2006
NEWS	16	FEB 28	MEDLINE/LMEDLINE reload improves functionality
NEWS	17	FEB 28	TOXCENTER reloaded with enhancements
NEWS	18	FEB 28	REGISTRY/ZREGISTRY enhanced with more experimental spectral property data
NEWS	19	MAR 01	INSPEC reloaded and enhanced
NEWS	20	MAR 03	Updates in PATDPA; addition of IPC 8 data without attributes
NEWS	21	MAR 08	X.25 communication option no longer available after June 2006
NEWS	22	MAR 22	EMBASE is now updated on a daily basis
NEWS	23	APR 03	New IPC 8 fields and IPC thesaurus added to PATDPAFULL
NEWS	24	APR 03	Bibliographic data updates resume; new IPC 8 fields and IPC thesaurus added in PCTFULL
NEWS	25	APR 04	STN AnaVist \$500 visualization usage credit offered
NEWS EXPRESS	FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005. V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT http://download.cas.org/express/v8.0-Discover/		
NEWS HOURS	STN Operating Hours Plus Help Desk Availability		
NEWS LOGIN	Welcome Banner and News Items		

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation

of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 12:20:43 ON 07 APR 2006

=> file medline embase biosis biotechds scisearch hcaplus ntis lifesci		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.42	0.42

FILE 'MEDLINE' ENTERED AT 12:21:52 ON 07 APR 2006

FILE 'EMBASE' ENTERED AT 12:21:52 ON 07 APR 2006
Copyright (c) 2006 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 12:21:52 ON 07 APR 2006
Copyright (c) 2006 The Thomson Corporation

FILE 'BIOTECHDS' ENTERED AT 12:21:52 ON 07 APR 2006
COPYRIGHT (C) 2006 THE THOMSON CORPORATION

FILE 'SCISEARCH' ENTERED AT 12:21:52 ON 07 APR 2006
Copyright (c) 2006 The Thomson Corporation

FILE 'HCAPLUS' ENTERED AT 12:21:52 ON 07 APR 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'NTIS' ENTERED AT 12:21:52 ON 07 APR 2006
Compiled and distributed by the NTIS, U.S. Department of Commerce.
It contains copyrighted material.
All rights reserved. (2006)

FILE 'LIFESCI' ENTERED AT 12:21:52 ON 07 APR 2006
COPYRIGHT (C) 2006 Cambridge Scientific Abstracts (CSA)

=> s "14171 kinase/"
<-----User Break----->

=> s "14171 kinase?"
L1 1 "14171 KINASE?"

=> d ibib ab

L1 ANSWER 1 OF 1 BIOTECHDS COPYRIGHT 2006 THE THOMSON CORP. on STN
ACCESSION NUMBER: 2004-12766 BIOTECHDS
TITLE: New 14171 protein kinase and nucleic acid, useful for
diagnosing or treating diseases with aberrant expression of
the 14171 protein kinase, such as cancer, an immunological
disorder, inflammation, heart failure and hypertension;
recombinant enzyme protein production via plasmid
expression in host cell for use in disease therapy
AUTHOR: KAPPELLER-LIBERMANN R
PATENT ASSIGNEE: MILLENNIUM PHARM INC
PATENT INFO: US 2004048305 11 Mar 2004
APPLICATION INFO: US 2003-658904 10 Sep 2003
PRIORITY INFO: US 2003-658904 10 Sep 2003; US 2000-182096 11 Feb 2000
DOCUMENT TYPE: Patent
LANGUAGE: English
OTHER SOURCE: WPI: 2004-226195 [21]

AB

DERWENT ABSTRACT:

NOVELTY - An isolated nucleic acid molecule (I) comprising a fully defined sequence of 3860 or 2355 base pairs (bp) (SEQ ID NO: 1 and 3) as given in the specification; a fragment of a fully defined sequence of 21 bp (SEQ ID NO: 21, 22 or 23) as given in the specification; or encoding a polypeptide having a fully defined sequence of 784 amino acids (SEQ ID NO: 2) as given in the specification, is new.

DETAILED DESCRIPTION - An isolated nucleic acid molecule comprises: (a) a fully defined sequence of 3860 or 2355 bp (SEQ ID NO: 1 and 3) as given in the specification; (b) a fragment of a fully defined sequence of 21 bp (SEQ ID NO: 21, 22 or 23) as given in the specification; (c) a nucleic acid molecule which encodes a polypeptide having a fully defined sequence of 784 amino acids (SEQ ID NO: 2) as given in the specification, or its fragment having at least 300 contiguous amino acids and kinase activity; or (d) the complement of (a), (b), (c), or (d). INDEPENDENT CLAIMS are also included for: (1) an expression construct comprising a recombinant nucleic acid molecule comprising the nucleic acid molecule (I); (2) a host cell comprising a recombinant nucleic acid molecule comprising the nucleic acid molecule (I); (3) an isolated polypeptide comprising: (a) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence with SEQ ID NO: 1 or 3; (b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, where the fragment comprises at least 300 contiguous amino acids of SEQ ID NO:2 and where at least 300 contiguous amino acids have kinase activity; (c) an antigenic fragment of SEQ ID NO:2 comprising at least 15 amino acid residues of SEQ ID NO:2; or (d) a polypeptide having the amino acid sequence of SEQ ID NO:2; (4) an antibody which selectively binds to a polypeptide of (3); (5) producing a polypeptide of (3), comprising culturing the host cell of (2) under conditions in which the nucleic acid molecule is expressed; (6) a kit comprising a compound which selectively binds to a polypeptide of (3) and instructions for use; (7) a kit comprising a compound which selectively hybridizes to a nucleic acid molecule (I) and instructions for use; (8) identifying a compound which binds to a polypeptide of (3), comprising contacting a polypeptide, or a cell expressing the polypeptide with a test compound and determining whether the polypeptide binds to the test compound; (9) modulating the activity of a polypeptide of (3), comprising contacting a polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide; (10) identifying a compound which modulates the activity of a polypeptide of (3), comprising contacting the polypeptide with a test compound and determining the effect of the test compound on the activity of the polypeptide to therefore identify a compound that modulates the activity of the polypeptide; (11) identifying a subject having a disorder or at risk of developing a disorder selected from the group consisting of cancer, an immunological disorder, a viral disorder and an apoptotic disorder, comprising contacting a sample obtained from the subject comprising nucleic acid molecules with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule (I), and detecting in the sample the presence of a nucleic acid molecule which hybridizes to the probe or primer, therefore identifying a subject having the disorder, or at risk for developing the disorder; or comprising contacting a sample obtained from the subject comprising polypeptides with a compound which selectively binds to the polypeptide of (3), and detecting in the sample the presence of a polypeptide which binds to the compound, therefore, identifying a subject having the disorder, or at risk for developing the disorder; and (12) treating a subject having a disorder selected from the group consisting of cancer, an immunological disorder, a viral disorder and an apoptotic disorder comprising administering to the subject an effective amount of an agent which targets the expression or activity of a nucleic acid molecule (I).

BIOTECHNOLOGY - Preferred Nucleic Acid: The nucleic acid further comprises nucleic acid sequences encoding a heterologous polypeptide. Preferred Polypeptide: The polypeptide of (3) further comprises

heterologous amino acid sequences. Preferred Antibody: The antibody preferably binds to an antigenic fragment of SEQ ID NO: 2 selected from the group consisting of a fully defined sequence of 21, 20 or 21 bp (base pairs) (SEQ ID NO: 17, 18 and 19), as given in the specification.

Preferred Method: The binding of the test compound to the polypeptide in the method of (8) is detected by detection of binding by direct detecting of test compound/polypeptide binding, detection of binding using a competition binding assay, or detection of binding using an assay for protein kinase-mediated phosphorylation. The activity of the polypeptide in the method of (10) is determined in a kinase assay using a 14171 kinase substrate. The nucleic acid probe or primer in the method of (11) is from a fully defined sequence of 20, 20 or 26 bp (SEQ ID NO: 9, 10 or 11) as given in the specification.

ACTIVITY - Cytostatic; Virucide; Antiinflammatory; Cardiant; Antiarrhythmic; Hypotensive. No biological data given.

MECHANISM OF ACTION - Protein-Kinase-Modulator. No biological data given.

USE - The methods and compositions of the present invention are useful for the diagnosis and/or treatment of diseases or conditions associated with aberrant expression or activity of a 14171 protein kinase, such as cancer, an immunological disorder, inflammation, heart failure, hypertension, atrial fibrillation, a viral disorder and an apoptotic disorder. They can also be used in chromosome mapping, tissue typing, predictive medicine, forensic biology and prognostic assays.

ADMINISTRATION - Dosage of the pharmaceutical composition ranges from 0.001-30 mg/kg body weight, preferably 5-6 mg/kg. Routes of administration of the pharmaceutical compositions include oral, pulmonary, intramuscular, intraperitoneal, intravenous, subcutaneous, inhalation, transdermal, nasal and rectal.

EXAMPLE - Total RNA was prepared from various human tissues by a single step extraction method using RNA STAT-60. Each RNA preparation was treated with DNase I at 37 degrees centigrade for 1 hour. DNase I treatment was determined to be complete if the sample required at least 38 PCR amplification cycles to reach a threshold level of fluorescence using beta-2 microglobulin as an internal amplicon reference. After phenol extraction cDNA was prepared from the sample using SUPERScript Choice System. A negative control of RNA without reverse transcriptase was mock reverse transcribed for each RNA sample. (62 pages)

=> d his

(FILE 'HOME' ENTERED AT 12:20:43 ON 07 APR 2006)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:21:52 ON 07 APR 2006

L1 1 S "14171 KINASE?"

=> s kinase?

L2 1434878 KINASE?

=> s clon? or express? or recombinant

L3 7629584 CLON? OR EXPRESS? OR RECOMBINANT

=> s "T-P" motif

L4 73 "T-P" MOTIF

=> s l2 and l4

L5 49 L2 AND L4

=> dup rem l5

PROCESSING COMPLETED FOR L5

L6 12 DUP REM L5 (37 DUPLICATES REMOVED)

=> d 1-12 ibib ab

L6 ANSWER 1 OF 12 MEDLINE on STN DUPLICATE 1
ACCESSION NUMBER: 2006100693 IN-PROCESS
DOCUMENT NUMBER: PubMed ID: 16365045
TITLE: The low density lipoprotein receptor-related protein 6
interacts with glycogen synthase kinase 3 and
attenuates activity.
AUTHOR: Mi Kaihong; Dolan Philip J; Johnson Gail V W
CORPORATE SOURCE: Department of Psychiatry, University of Alabama at
Birmingham, Birmingham, Alabama 35294-0017, USA.
CONTRACT NUMBER: NS051279 (NINDS)
SOURCE: The Journal of biological chemistry, (2006 Feb 24) Vol.
281, No. 8, pp. 4787-94. Electronic Publication:
2005-12-19.
Journal code: 2985121R. ISSN: 0021-9258.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: NONMEDLINE; IN-PROCESS; NONINDEXED; Priority Journals
ENTRY DATE: Entered STN: 20060222
Last Updated on STN: 20060314
AB Glycogen synthase kinase 3 (GSK3) is a widely expressed Ser/Thr
protein kinase that phosphorylates numerous substrates. This
large number of substrates requires precise and specific regulation of
GSK3 activity, which is achieved by a combination of phosphorylation,
localization, and interactions with GSK3-binding proteins. Members of the
Wnt canonical pathway have been shown to influence GSK3 activity. Through
a yeast two-hybrid screen, we identified the Wnt canonical pathway
co-receptor protein low density lipoprotein receptor-related protein 6
(LRP6) as a GSK3-binding protein. The interaction between the C terminus
of LRP6 and GSK3 was also confirmed by in vitro GST pull-down assays and
in situ coimmunoprecipitation assays. In vitro assays using
immunoprecipitated proteins demonstrated that the C terminus of LRP6
significantly attenuated the activity of GSK3beta. In situ, LRP6
significantly decreased GSK3beta-mediated phosphorylation of tau at both
primed and unprimed sites. Finally, it was also demonstrated that
GSK3beta phosphorylates the PPP(S/T)P motifs
in the C terminus of LRP6. This is the first identification of a direct
interaction between LRP6 and GSK3, which results in an attenuation of GSK3
activity.

L6 ANSWER 2 OF 12 MEDLINE on STN DUPLICATE 2
ACCESSION NUMBER: 2004100692 MEDLINE
DOCUMENT NUMBER: PubMed ID: 14681225
TITLE: Extracellular signal-regulated kinases 1/2 are
serum-stimulated "Bim(EL) kinases" that bind to
the BH3-only protein Bim(EL) causing its phosphorylation
and turnover.
AUTHOR: Ley Rebecca; Ewings Katherine E; Hadfield Kathryn; Howes
Elizabeth; Balmano Kathryn; Cook Simon J
CORPORATE SOURCE: Laboratory of Molecular Signalling, Signalling Programme,
The Babraham Institute, Cambridge CB2 4AT, United Kingdom..
becky.ley@bbsrc.ac.uk
SOURCE: The Journal of biological chemistry, (2004 Mar 5) Vol. 279,
No. 10, pp. 8837-47. Electronic Publication: 2003-12-17.
Journal code: 2985121R. ISSN: 0021-9258.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200407
ENTRY DATE: Entered STN: 20040302
Last Updated on STN: 20040707

Entered Medline: 20040706

AB Bim, a "BH3-only" protein, is expressed de novo following withdrawal of serum survival factors and promotes cell death. We have shown previously that activation of the ERK1/2 pathway promotes phosphorylation of Bim(EL), targeting it for degradation via the proteasome. However, the nature of the kinase responsible for Bim(EL) phosphorylation remained unclear. We now show that Bim(EL) is phosphorylated on at least three sites in response to activation of the ERK1/2 pathway. By using the peptidylprolyl isomerase, Pin1, as a probe for proline-directed phosphorylation, we show that ERK1/2-dependent phosphorylation of Bim(EL) occurs at (S/T)P motifs. ERK1/2 phosphorylates Bim(EL), but not Bim(S) or Bim(L), in vitro, and mutation of Ser(65) to alanine blocks the phosphorylation of Bim(EL) by ERK1/2 in vitro and in vivo and prevents the degradation of the protein following activation of the ERK1/2 pathway. We also find that ERK1/2, but not JNK, can physically associate with GST-Bim(EL), but not GST-Bim(L) or GST-Bim(S), in vitro. ERK1/2 also binds to full-length Bim(EL) in vivo, and we have localized a potential ERK1/2 "docking domain" lying within a 27-amino acid stretch of the Bim(EL) protein. Our findings provide new insights into the post-translational regulation of Bim(EL) and the role of the ERK1/2 pathway in cell survival signaling.

L6 ANSWER 3 OF 12 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
DUPLICATE 3

ACCESSION NUMBER: 2005:89678 BIOSIS

DOCUMENT NUMBER: PREV200500087142

TITLE: Physiological role of the oxidative stress-susceptible
TRPM2 Ca2+ channel in immunocytes.

AUTHOR(S): Yamamoto, Shinichiro [Reprint Author]; Shimizu, Shunichi;
Ishii, Masakazu; Hagiwara, Tamio; Hara, Yuji; Negoro,
Takaharu; Nishida, Motohiro; Tobe, Takashi; Mori, Yasuo;
Kiuchi, Yuji

CORPORATE SOURCE: Grad Sch EngnDept Synthet Chem and BiolMol Biol Lab, Kyoto
Univ, Kyoto, 6068501, Japan

SOURCE: Yakugaku Zasshi, (2004) Vol. 124, No. Suppl. 4, pp.
237-240. print.

ISSN: 0031-6903 (ISSN print).

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 2 Mar 2005

Last Updated on STN: 2 Mar 2005

AB TRPM2 is a Ca2+ permeable channel activated by various triggers including the oxidative stress including hydrogen peroxide (H2O2). TRPM2 is expressed in immunocytes such as monocytes, lymphocytes, and neutrophils. However its physiological role is unclear. Although the activation of TRPM2 by H2O2 seems to be mediated by NAD+ and/or ADP-ribose, the activation mechanisms in the context of physiological signaling are not elucidated. Thus, We investigated the activation mechanisms of TRPM2 and the physiological role of Ca2+ influx via TRPM2 using monocytic cell line U937. Addition of H2O2 to U937 cells triggered Ca2+ influx, and the both Ca2+ influx and TRPM2 expression were attenuated by the treatment with TRPM2-specific siRNA. The H2O2-triggered TRPM2 activation was also inhibited by the treatment with ERK kinase inhibitor, PD98059. Moreover, the activation of TRPM2 recombinantly expressed in HEK293 cells was blocked by the mutation of putative phosphorylation sites (S/T -P motif) by ERK, suggesting that H2O2-triggered TRPM2 activation was controlled by ERK pathway. In U937 cells, H2O2 induced interleukin-8 (IL-8) production in extracellular Ca2+ dependent manner, which was inhibited by the treatment with TRPM2 specific siRNA and PD98059. The Ca2+ influx via TRPM2 induced by H2O2 participates in IL-8 production in U937 cells.

L6 ANSWER 4 OF 12 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights
reserved on STN

ACCESSION NUMBER: 2004000668 EMBASE
 TITLE: [Unexpected roles of the peptidyl-prolyl cis/trans isomerase Pin1].
 PIN1: UNE PEPTIDYL-PROLYL CIS/TRANS ISOMERASE AUX ROLES INSOUPECONNES.
 AUTHOR: Lavoie S.B.; Albert A.L.; Vincent M.
 CORPORATE SOURCE: S.B. Lavoie, Departement de Medecine et CREFSIP, Pavillon C.E. Marchand, Universite Laval, Quebec, Que. G1K 7P4, Canada. seb_lavoie@iquebec.com
 SOURCE: Medecine/Sciences, (2003) Vol. 19, No. 12, pp. 1251-1258. .
 Refs: 40
 ISSN: 0767-0974 CODEN: MSMSE4
 COUNTRY: France
 DOCUMENT TYPE: Journal; General Review
 FILE SEGMENT: 005 General Pathology and Pathological Anatomy
 008 Neurology and Neurosurgery
 029 Clinical Biochemistry
 LANGUAGE: French
 SUMMARY LANGUAGE: English; French
 ENTRY DATE: Entered STN: 16 Jan 2004
 Last Updated on STN: 16 Jan 2004

AB Peptidyl-prolyl isomerases (PPlases) are chaperone enzymes which alter the peptide bond between a given amino acid and a proline, changing it from the as to the trans conformation and vice versa. This modification can cause dramatic structural modifications which can affect the properties of targeted proteins. The ubiquitous PPlase Pin1, conserved from yeast to human, has been shown to be necessary for entry into mitosis. The yeast homologue, Ess1, is essential for cell survival. Pin1 possesses a WW domain which specifically recognizes pSer-Pro and pThr-Pro motifs in which the first amino acid is phosphorylated. Pin1 binds to many proteins implicated in cell cycle regulation (e.g. p53, Myt1, Wee1, and Cdc25C). Pin1 also targets tau, a protein forming part of the neuronal cytoskeleton which is hyperphosphorylated in patients suffering from Alzheimer's disease (AD). Pin1 could, therefore, be involved in the pathogenesis of AD. Furthermore, Pin1 also binds two proteins involved in transcription: Rpb1, the largest subunit of RNA polymerase II and Spt5, a regulator of the elongation of transcription. Both these proteins possess domains rich in S/T-P motifs which can be targeted by Pin1 when phosphorylated. Recent studies show that Pin1 modulates the dephosphorylation of some proteins by allowing trans-specific phosphatases to recognize their target after isomerization. This unexpected role might allow protein regulation via peptidyl-prolyl isomerase activity.

L6 ANSWER 5 OF 12 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
 ACCESSION NUMBER: 2001:245224 BIOSIS
 DOCUMENT NUMBER: PREV200100245224
 TITLE: IRS-1-dependent IFNalpha signaling is impaired by a FRAP regulated mechanism.
 AUTHOR(S): Hartman, Matthew E. [Reprint author]; Villela-Bach, Montserrat [Reprint author]; Chen, Jie; Freund, Gregory G. [Reprint author]
 CORPORATE SOURCE: University of Illinois at Urbana-Champaign, 1201 West Gregory, 261 ERLM, Urbana, IL, 61801, USA
 SOURCE: FASEB Journal, (March 8, 2001) Vol. 15, No. 5, pp. A945. print.
 Meeting Info.: Annual Meeting of the Federation of American Societies for Experimental Biology on Experimental Biology 2001. Orlando, Florida, USA. March 31-April 04, 2001.
 CODEN: FAJOEC. ISSN: 0892-6638.
 DOCUMENT TYPE: Conference; (Meeting)
 Conference; Abstract; (Meeting Abstract)
 LANGUAGE: English
 ENTRY DATE: Entered STN: 23 May 2001
 Last Updated on STN: 19 Feb 2002

AB We have previously shown that interferon-alpha (IFNalpha)-dependent tyrosine phosphorylation of insulin receptor substrate-1 (IRS-1) and subsequent IRS-1 phosphatidylinositol 3-kinase (PI 3-kinase) association/activation is impaired by serine phosphorylation of IRS-1 due to the reduced ability of serine phosphorylated IRS-1 to serve as a substrate for Janus kinase 1 (JAK1). Here we report that FKBP12-rapamycin associated protein (FRAP) is a physiologic IRS-1 serine kinase that blocks IFNalpha signaling by serine phosphorylating IRS-1. We found that treatment of U266 cells with the FRAP inhibitor rapamycin increased IFNalpha-dependent tyrosine phosphorylation by 2-fold while reducing constitutive IRS-1 serine phosphorylation within S/T-P motifs by 80%. On the contrary, wortmannin treatment had no effect on IFNalpha stimulated IRS-1 tyrosine phosphorylation. Importantly, both FRAP and insulin-activated p70s6k, serine phosphorylated IRS-1 between residues 511-772 (IRS-1511-772), but only FRAP-dependent IRS-1511-772 serine phosphorylation inhibited by 50% subsequent JAK1-dependent tyrosine phosphorylation of IRS-1. Taken together, these data indicate that FRAP, but not p70s6k, is an in vivo IRS-1 serine kinase that negatively regulates JAK1-dependent IRS-1 tyrosine phosphorylation and suggest that FRAP may modulate cytokine signal transduction through IRSs.

L6 ANSWER 6 OF 12 MEDLINE on STN DUPLICATE 4
 ACCESSION NUMBER: 2001179452 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 11162588
 TITLE: Frap-dependent serine phosphorylation of IRS-1 inhibits IRS-1 tyrosine phosphorylation.
 AUTHOR: Hartman M E; Villela-Bach M; Chen J; Freund G G
 CORPORATE SOURCE: Department of Animal Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA.
 CONTRACT NUMBER: CA-61931 (NCI)
 GM-58064 (NIGMS)
 SOURCE: Biochemical and biophysical research communications, (2001 Jan 26) Vol. 280, No. 3, pp. 776-81.
 Journal code: 0372516. ISSN: 0006-291X.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200103
 ENTRY DATE: Entered STN: 20010404
 Last Updated on STN: 20010404
 Entered Medline: 20010329

AB We have previously shown that interferon-alpha (IFN alpha)-dependent tyrosine phosphorylation of insulin receptor substrate-1 (IRS-1) is impaired by serine phosphorylation of IRS-1 due to the reduced ability of serine phosphorylated IRS-1 to serve as a substrate for Janus kinase 1 (JAK1). Here we report that FKBP12-rapamycin-associated protein (FRAP) is a physiologic IRS-1 kinase that blocks IFN alpha signaling by serine phosphorylating IRS-1. We found that both FRAP and insulin-activated p70 S6 kinase (p70(s6k)) serine phosphorylated IRS-1 between residues 511 and 772 (IRS-1(511-772)). Importantly, only FRAP-dependent IRS-1(511-772) serine phosphorylation inhibited by 50% subsequent JAK1-dependent tyrosine phosphorylation of IRS-1. Furthermore, treatment of U266 cells with the FRAP inhibitor rapamycin increased IFN alpha-dependent tyrosine phosphorylation by twofold while reducing constitutive IRS-1 serine phosphorylation within S/T-P motifs by 80%. Taken together, these data indicate that FRAP, but not p70(s6k), is a likely physiologic IRS-1 serine kinase that negatively regulates JAK1-dependent IRS-1 tyrosine phosphorylation and suggests that FRAP may modulate IRS-dependent cytokine signaling.
 Copyright 2001 Academic Press.

L6 ANSWER 7 OF 12 MEDLINE on STN DUPLICATE 5
 ACCESSION NUMBER: 2000105769 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 10637505
 TITLE: ERK activation induces phosphorylation of Elk-1 at multiple S/T-P motifs to high stoichiometry.
 AUTHOR: Cruzalegui F H; Cano E; Treisman R
 CORPORATE SOURCE: Transcription Laboratory, Imperial Cancer Research Fund, 44 Lincoln's Inn Fields, London WC2A 3PX, UK.
 SOURCE: Oncogene, (1999 Dec 23) Vol. 18, No. 56, pp. 7948-57. Journal code: 8711562. ISSN: 0950-9232.
 PUB. COUNTRY: ENGLAND: United Kingdom
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200002
 ENTRY DATE: Entered STN: 20000218
 Last Updated on STN: 20000218
 Entered Medline: 20000204

AB Elk-1, a member of the TCF family of Ets domain proteins, contains a C-terminal transcriptional activation domain with multiple copies of the MAPK core consensus sequence S/T-P. This region is phosphorylated by MAP kinases in vitro and in vivo, but the extent and kinetics of phosphorylation at the different sites have not been investigated in detail. We prepared antisera against the phosphorylated forms of residues T353, T363, T368, S383, S389 and T417. The antisera specifically recognize the phosphorylated Elk-1 C terminus and are specific for their cognate sites, as assessed by peptide competition and mutagenesis experiments. Analysis of cells stably expressing Elk-1 in vivo shows that following serum or TPA stimulation, residues T353, T363, T368, S383, S389 and T417 become phosphorylated with similar kinetics. Mutation of any one site does not prevent phosphorylation of the others. Mutation to alanine of S383, F378 or W379, which virtually abolishes transcriptional activation by Elk-1, does not affect phosphorylation of any sites tested. Analysis of Elk-1 using two-dimensional gel electrophoresis shows that following ERK activation Elk-1 receives at least six phosphates in addition to those present prior to stimulation. We propose that the Elk-1 C-terminal regulatory domain becomes stoichiometrically phosphorylated following growth factor stimulation.

L6 ANSWER 8 OF 12 MEDLINE on STN DUPLICATE 6
 ACCESSION NUMBER: 1999009106 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 9792705
 TITLE: Mitogen-activated protein kinase phosphorylates and regulates the HIV-1 Vif protein.
 AUTHOR: Yang X; Gabuzda D
 CORPORATE SOURCE: Department of Cancer Immunology & AIDS, Dana-Farber Cancer Institute, Boston, Massachusetts 02115, USA.
 CONTRACT NUMBER: AI28691 (NIAID)
 AI36186 (NIAID)
 AO6514
 +
 SOURCE: The Journal of biological chemistry, (1998 Nov 6) Vol. 273, No. 45, pp. 29879-87. Journal code: 2985121R. ISSN: 0021-9258.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals; AIDS
 ENTRY MONTH: 199812
 ENTRY DATE: Entered STN: 19990115
 Last Updated on STN: 19990115
 Entered Medline: 19981210

AB The human immunodeficiency virus type 1 (HIV-1) Vif protein plays a

critical role in virus replication and infectivity. Here we show that Vif is phosphorylated and regulated by p44/42 mitogen-activated protein kinase (MAPK). Vif phosphorylation by MAPK was demonstrated in vitro as well as in vivo and was shown to occur on serine and threonine residues. Two-dimensional tryptic phosphopeptide mapping indicated that Vif is phosphorylated by MAPK on the same sites in vitro and in vivo. Radioactive peptide sequencing identified two phosphorylation sites, Thr96 and Ser165. These phosphorylation sites do not correspond to the known optimum consensus sequences for phosphorylation by MAPK (PX(S/T)P) nor to the minimum consensus sequence ((S/T)P), indicating that MAPK can phosphorylate proteins at sites other than those containing the PX(S/T)P or (S/T)P motifs. Synthetic Vif peptides corresponding to the local sequences of the phosphorylation sites were not phosphorylated by MAPK, suggesting that recognition of these sites by MAPK is likely to require structural determinants outside the phosphorylation site. Mutations of the Thr96 site, which is conserved among Vif sequences from HIV-1, HIV-2, and SIV, resulted in significant loss of Vif activity and inhibition of HIV-1 replication. These results suggest that MAPK plays a direct role in regulating HIV-1 replication and infectivity by phosphorylating Vif and identify a novel mechanism for activation of HIV-1 replication by mitogens and other extracellular stimuli.

L6 ANSWER 9 OF 12 MEDLINE on STN DUPLICATE 7
 ACCESSION NUMBER: 1999030926 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 9811754
 TITLE: Host cell-virus cross talk: phosphorylation of a hepatitis B virus envelope protein mediates intracellular signaling.
 AUTHOR: Rothmann K; Schnolzer M; Radziwill G; Hildt E; Moelling K; Schaller H
 CORPORATE SOURCE: Zentrum fur Molekulare Biologie Heidelberg, D-69124 Heidelberg, Germany.
 SOURCE: Journal of virology, (1998 Dec) Vol. 72, No. 12, pp. 10138-47.
 Journal code: 0113724. ISSN: 0022-538X.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199811
 ENTRY DATE: Entered STN: 19990115
 Last Updated on STN: 20000303
 Entered Medline: 19981130

AB Phosphorylation of cytosolic pre-S domains of the duck hepatitis B virus (DHBV) large envelope protein (L) was identified as a regulatory modification involved in intracellular signaling. By using biochemical and mass spectrometric analyses of phosphopeptides obtained from metabolically radiolabeled L protein, a single phosphorylation site was identified at serine 118 as part of a PX(S/T)P motif, which is strongly preferred by ERK-type mitogen-activated protein kinases (MAP kinases). ERK2 specifically phosphorylated L at serine 118 in vitro, and L phosphorylation was inhibited by a coexpressed MAP kinase-specific phosphatase. Furthermore, L phosphorylation and ERK activation were shown to be induced in parallel by various stimuli. Functional analysis with transfected cells showed that DHBV L possesses the ability to activate gene expression in trans and, by using mutations eliminating (S-->A) or mimicking (S-->D) serine phosphorylation, that this function correlates with L phosphorylation. These mutations had, however, no major effects on virus production in cell culture and in vivo, indicating that L phosphorylation and transactivation are not essential for hepadnavirus replication and morphogenesis. Together, these data suggest a role of the L protein in intracellular host-virus cross talk by varying the levels of pre-S phosphorylation in response to the state of the cell.

L6 ANSWER 10 OF 12 MEDLINE on STN DUPLICATE 8
 ACCESSION NUMBER: 1998256424 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 9592139
 TITLE: SCP2: a major protein component of the axial elements of synaptonemal complexes of the rat.
 AUTHOR: Offenberg H H; Schalk J A; Meuwissen R L; van Aalderen M; Kester H A; Dietrich A J; Heyting C
 CORPORATE SOURCE: Department of Genetics, Agricultural University, Dreijenlaan 2, NL-6703 HA Wageningen, The Netherlands.
 SOURCE: Nucleic acids research, (1998 Jun 1) Vol. 26, No. 11, pp. 2572-9.
 Journal code: 0411011. ISSN: 0305-1048.
 PUB. COUNTRY: ENGLAND: United Kingdom
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 OTHER SOURCE: GENBANK-Y08981; GENBANK-Y08982
 ENTRY MONTH: 199807
 ENTRY DATE: Entered STN: 19980731
 Last Updated on STN: 20000303
 Entered Medline: 19980720

AB In the axial elements of synaptonemal complexes (SCs) of the rat, major protein components have been identified, with relative electrophoretic mobilities (M_{rs}) of 30 000-33 000 and 190 000. Using monoclonal anti-SC antibodies, we isolated cDNA fragments which encode the 190 000 M_r component of rat SCs. The translation product predicted from the nucleotide sequence of the cDNA, called SCP2 (for synaptonemal complex protein 2), is a basic protein (pI = 8.0) with a molecular mass of 173 kDa. At the C-terminus, a stretch of approximately 50 amino acid residues is predicted to be capable of forming coiled-coil structures. SCP2 contains two clusters of S/T-P motifs, which are common in DNA-binding proteins. These clusters flank the central, most basic part of the protein (pI = 9.5). Three of the S/T-P motifs are potential target sites for p34(cdc2) protein kinase. In addition, SCP2 has eight potential CAMP/cGMP-dependent protein kinase target sites. The gene encoding SCP2 is transcribed specifically in the testis, in meiotic prophase cells. At the amino acid sequence and secondary structural level, SCP2 shows some similarity to the Red1 protein, which is involved in meiotic recombination and the assembly of axial elements of SCs in yeast. We speculate that SCP2 is a DNA-binding protein involved in the structural organization of meiotic prophase chromosomes.

L6 ANSWER 11 OF 12 SCISEARCH COPYRIGHT (c) 2006 The Thomson Corporation on STN DUPLICATE 9
 ACCESSION NUMBER: 1995:391391 SCISEARCH
 THE GENUINE ARTICLE: RC664
 TITLE: COMPARATIVE-ANALYSIS OF THE TERNARY COMPLEX FACTORS ELK-1, SAP-1A AND SAP-2 (ERP/NET)
 AUTHOR: PRICE M A (Reprint); ROGERS A E; TREISMAN R
 CORPORATE SOURCE: IMPERIAL CANC RES FUND, TRANSCRIPT LAB, 44 LINCOLNS INN FIELDS, LONDON WC2A 3PX, ENGLAND (Reprint)
 COUNTRY OF AUTHOR: ENGLAND
 SOURCE: EMBO JOURNAL, (1 JUN 1995) Vol. 14, No. 11, pp. 2589-2601. ISSN: 0261-4189.
 PUBLISHER: OXFORD UNIV PRESS UNITED KINGDOM, WALTON ST JOURNALS DEPT, OXFORD, ENGLAND OX2 6DP.
 DOCUMENT TYPE: Article; Journal
 FILE SEGMENT: LIFE
 LANGUAGE: English
 REFERENCE COUNT: 45
 ENTRY DATE: Entered STN: 1995
 Last Updated on STN: 1995
 ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS

AB A transcription factor ternary complex composed of Serum Response Factor (SRF) and Ternary Complex Factor (TCF) mediates the response of the c-fos Serum Response Element (SRE) to growth factors and mitogens. Three Ets domain proteins, Elk-1, SAP-1 and ERP/NET, have been reported to have the properties of TCF. Here we compare Elk-1 and SAP-1a with the human ERP/NET homologue SAP-2. All three TCF RNAs are ubiquitously expressed at similar relative levels. All three proteins contain conserved regions that interact with SRF and the c-fos SRE with comparable efficiency, but in vitro complex formation by SAP-2 is strongly inhibited by its C-terminal sequences. Similarly, only Elk-1 and SAP-1a efficiently bind the c-fos SRE in vivo; ternary complex formation by SAP-2 is weak and is substantially unaffected by serum stimulation or v-ras co-expression. All three TCFs contain C-terminal transcriptional activation domains that are phosphorylated following growth factor stimulation. Activation requires conserved S/T-P motifs found in all the TCF family members, Each TCF activation domain can be phosphorylated in vitro by partially purified ERK2, and ERK activation in vivo is sufficient to potentiate transcriptional activation.

L6 ANSWER 12 OF 12 MEDLINE on STN DUPLICATE 10
 ACCESSION NUMBER: 92317057 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 1618840
 TITLE: Purification and characterization of a novel proline-directed protein kinase from bovine brain.
 AUTHOR: Lew J; Beaudette K; Litwin C M; Wang J H
 CORPORATE SOURCE: Department of Medical Biochemistry, Faculty of Medicine, University of Calgary, Alberta, Canada.
 SOURCE: The Journal of biological chemistry, (1992 Jul 5) Vol. 267, No. 19, pp. 13383-90.
 Journal code: 2985121R. ISSN: 0021-9258.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199208
 ENTRY DATE: Entered STN: 19920815
 Last Updated on STN: 19970203
 Entered Medline: 19920805

AB A novel protein kinase which phosphorylates a synthetic peptide substrate (RRPDAHRTPNRAF) has been purified approximately 200,000-fold from bovine brain. This peptide contains the consensus sequence for phosphorylation by the p34cdc2 kinase. The purification procedure took advantage of the phenomenon that this novel brain kinase, in partially purified extracts, chromatographed on a gel filtration column as a high molecular weight complex which dissociated in buffer containing 1 M NaCl. The purified native enzyme was estimated to be approximately 63,000, and displayed two bands of M(r) = 33,000 and 25,000 on sodium dodecyl sulfate-polyacrylamide gel electrophoresis. On Western immunoblot, the M(r) = 33,000 peptide reacted strongly with antibodies specific for a conserved amino-terminal sequence, weakly with antibodies to the conserved PSTAIRE sequence, and not at all with antibodies to the carboxyl terminus, of HeLa cell p34cdc2. The brain kinase and p34cdc2 were similar in displaying good activity toward the parent peptide substrate, but no activity toward peptide analogues in which the -T-P- motif was substituted with either -T-G- or -T-A-. Both kinases showed marked preference in phosphorylating a peptide derived from H1 histone (KTPKKAKKPKTPKKAKKL), and both kinases could be phosphorylated by the src-family tyrosine kinase, p56lyn, purified from bovine spleen. However, the brain kinase did not co-purify with a subunit having a molecular weight corresponding to known cyclins, nor did it undergo specific interaction with p13suc1 beads, suggesting that this enzyme is distinct from p34cdc2.

=> d his

(FILE 'HOME' ENTERED AT 12:20:43 ON 07 APR 2006)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:21:52 ON 07 APR 2006

L1 1 S "14171 KINASE?"
L2 1434878 S KINASE?
L3 7629584 S CLON? OR EXPRESS? OR RECOMBINANT
L4 73 S "T-P" MOTIF
L5 49 S L2 AND L4
L6 12 DUP REM L5 (37 DUPLICATES REMOVED)

=> s (modulat? or activat? or inhibit?) and l2
L7 947418 (MODULAT? OR ACTIVAT? OR INHIBIT?) AND L2

=> s "14171"
L8 42 "14171"

=> s l7 and l8
L9 2 L7 AND L8

=> d 1-2 ibib ab

L9 ANSWER 1 OF 2 BIOTECHDS COPYRIGHT 2006 THE THOMSON CORP. on STN
ACCESSION NUMBER: 2004-12766 BIOTECHDS

TITLE: New 14171 protein **kinase** and nucleic acid, useful for diagnosing or treating diseases with aberrant expression of the 14171 protein **kinase**, such as cancer, an immunological disorder, inflammation, heart failure and hypertension; recombinant enzyme protein production via plasmid expression in host cell for use in disease therapy

AUTHOR: KAPELLER-LIBERMANN R
PATENT ASSIGNEE: MILLENNIUM PHARM INC
PATENT INFO: US 2004048305 11 Mar 2004
APPLICATION INFO: US 2003-658904 10 Sep 2003
PRIORITY INFO: US 2003-658904 10 Sep 2003; US 2000-182096 11 Feb 2000
DOCUMENT TYPE: Patent
LANGUAGE: English
OTHER SOURCE: WPI: 2004-226195 [21]

AB DERWENT ABSTRACT:

NOVELTY - An isolated nucleic acid molecule (I) comprising a fully defined sequence of 3860 or 2355 base pairs (bp) (SEQ ID NO: 1 and 3) as given in the specification; a fragment of a fully defined sequence of 21 bp (SEQ ID NO: 21, 22 or 23) as given in the specification; or encoding a polypeptide having a fully defined sequence of 784 amino acids (SEQ ID NO: 2) as given in the specification, is new.

DETAILED DESCRIPTION - An isolated nucleic acid molecule comprises: (a) a fully defined sequence of 3860 or 2355 bp (SEQ ID NO: 1 and 3) as given in the specification; (b) a fragment of a fully defined sequence of 21 bp (SEQ ID NO: 21, 22 or 23) as given in the specification; (c) a nucleic acid molecule which encodes a polypeptide having a fully defined sequence of 784 amino acids (SEQ ID NO: 2) as given in the specification, or its fragment having at least 300 contiguous amino acids and **kinase** activity; or (d) the complement of (a), (b), (c), or (d).

INDEPENDENT CLAIMS are also included for: (1) an expression construct comprising a recombinant nucleic acid molecule comprising the nucleic acid molecule (I); (2) a host cell comprising a recombinant nucleic acid molecule comprising the nucleic acid molecule (I); (3) an isolated polypeptide comprising: (a) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence with SEQ ID NO: 1 or 3;

(b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, where the fragment comprises at least 300 contiguous amino acids of SEQ ID NO:2 and where at least 300 contiguous amino acids have kinase activity; (c) an antigenic fragment of SEQ ID NO:2 comprising at least 15 amino acid residues of SEQ ID NO:2; or (d) a polypeptide having the amino acid sequence of SEQ ID NO:2; (4) an antibody which selectively binds to a polypeptide of (3); (5) producing a polypeptide of (3), comprising culturing the host cell of (2) under conditions in which the nucleic acid molecule is expressed; (6) a kit comprising a compound which selectively binds to a polypeptide of (3) and instructions for use; (7) a kit comprising a compound which selectively hybridizes to a nucleic acid molecule (I) and instructions for use; (8) identifying a compound which binds to a polypeptide of (3), comprising contacting a polypeptide, or a cell expressing the polypeptide with a test compound and determining whether the polypeptide binds to the test compound; (9) modulating the activity of a polypeptide of (3), comprising contacting a polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide; (10) identifying a compound which modulates the activity of a polypeptide of (3), comprising contacting the polypeptide with a test compound and determining the effect of the test compound on the activity of the polypeptide to therefore identify a compound that modulates the activity of the polypeptide; (11) identifying a subject having a disorder or at risk of developing a disorder selected from the group consisting of cancer, an immunological disorder, a viral disorder and an apoptotic disorder, comprising contacting a sample obtained from the subject comprising nucleic acid molecules with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule (I), and detecting in the sample the presence of a nucleic acid molecule which hybridizes to the probe or primer, therefore identifying a subject having the disorder, or at risk for developing the disorder; or comprising contacting a sample obtained from the subject comprising polypeptides with a compound which selectively binds to the polypeptide of (3), and detecting in the sample the presence of a polypeptide which binds to the compound, therefore, identifying a subject having the disorder, or at risk for developing the disorder; and (12) treating a subject having a disorder selected from the group consisting of cancer, an immunological disorder, a viral disorder and an apoptotic disorder comprising administering to the subject an effective amount of an agent which targets the expression or activity of a nucleic acid molecule (I).

BIOTECHNOLOGY - Preferred Nucleic Acid: The nucleic acid further comprises nucleic acid sequences encoding a heterologous polypeptide. **Preferred Polypeptide:** The polypeptide of (3) further comprises heterologous amino acid sequences. **Preferred Antibody:** The antibody preferably binds to an antigenic fragment of SEQ ID NO: 2 selected from the group consisting of a fully defined sequence of 21, 20 or 21 bp (base pairs) (SEQ ID NO: 17, 18 and 19), as given in the specification. **Preferred Method:** The binding of the test compound to the polypeptide in the method of (8) is detected by detection of binding by direct detecting of test compound/polypeptide binding, detection of binding using a competition binding assay, or detection of binding using an assay for protein kinase-mediated phosphorylation. The activity of the polypeptide in the method of (10) is determined in a kinase assay using a 14171 kinase substrate. The nucleic acid probe or primer in the method of (11) is from a fully defined sequence of 20, 20 or 26 bp (SEQ ID NO: 9, 10 or 11) as given in the specification.

ACTIVITY - Cytostatic; Virucide; Antiinflammatory; Cardiant; Antiarrhythmic; Hypotensive. No biological data given.

MECHANISM OF ACTION - Protein-Kinase-Modulator.
No biological data given.

USE - The methods and compositions of the present invention are

useful for the diagnosis and/or treatment of diseases or conditions associated with aberrant expression or activity of a 14171 protein kinase, such as cancer, an immunological disorder, inflammation, heart failure, hypertension, atrial fibrillation, a viral disorder and an apoptotic disorder. They can also be used in chromosome mapping, tissue typing, predictive medicine, forensic biology and prognostic assays.

ADMINISTRATION - Dosage of the pharmaceutical composition ranges from 0.001-30 mg/kg body weight, preferably 5-6 mg/kg. Routes of administration of the pharmaceutical compositions include oral, pulmonary, intramuscular, intraperitoneal, intravenous, subcutaneous, inhalation, transdermal, nasal and rectal.

EXAMPLE - Total RNA was prepared from various human tissues by a single step extraction method using RNA STAT-60. Each RNA preparation was treated with DNase I at 37 degrees centigrade for 1 hour. DNase I treatment was determined to be complete if the sample required at least 38 PCR amplification cycles to reach a threshold level of fluorescence using beta-2 microglobulin as an internal amplicon reference. After phenol extraction cDNA was prepared from the sample using SUPERScript Choice System. A negative control of RNA without reverse transcriptase was mock reverse transcribed for each RNA sample. (62 pages)

L9 ANSWER 2 OF 2 BIOTECHDS COPYRIGHT 2006 THE THOMSON CORP. on STN

ACCESSION NUMBER: 2003-28808 BIOTECHDS

TITLE: New 14171 human protein kinase and nucleic acids encoding the protein, useful for treating viral infections, cellular growth related disorders, cancers, disorders related with programmed cell death, or autoimmune disorders;
vector-mediated protein-kinase gene transfer and expression in host cell for recombinant protein production, drug screening and gene therapy

AUTHOR: KAPPELLER-LIBERMANN R

PATENT ASSIGNEE: MILLENNIUM PHARM INC

PATENT INFO: US 6630335 7 Oct 2003

APPLICATION INFO: US 2001-781882 12 Feb 2001

PRIORITY INFO: US 2001-781882 12 Feb 2001; US 2000-182096 11 Feb 2000

DOCUMENT TYPE: Patent

LANGUAGE: English

OTHER SOURCE: WPI: 2003-810551 [76]

AB DERWENT ABSTRACT:

NOVELTY - An isolated nucleic acid molecule (I) comprising: (a) a sequence of 3860 or 2355 bp given in the specification, or its complement; or (b) a sequence which encodes a polypeptide comprising a sequence of 784 amino acids (II) or the sequence (II) having a substitution for aspartate at position 143, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for: (1) a vector comprising (I); (2) a host cell comprising the vector; and (3) a method of producing a polypeptide comprising culturing the host cell of (2) under conditions in which the nucleic acid molecule is expressed to produce the polypeptide.

WIDER DISCLOSURE - (1) antibodies that selectively bind protein kinase polypeptide and fragments; (2) a method for detecting protein kinase activity of expression in a biological sample; (3) a method for modulating protein kinase activity; (4) a diagnostic assay for identifying the presence or absence of a genetic lesion for mutation characterized by aberrant modification or mutation of a gene encoding a protein kinase, misregulation of a gene encoding a protein kinase, or aberrant post-translational modification of a protein kinase; (5) a method for identifying a compound that binds to or modulates protein kinase activity; (6) a method for identifying compound that modulates the expression of a protein kinase gene; and (7) compound identified by the screening methods.

BIOTECHNOLOGY - Preferred Nucleic Acid: (I) further comprises nucleic acid sequences encoding a heterologous polypeptide. (I) comprises a sequence encoding a polypeptide comprising (II). Preferred Vector: The vector of comprises a nucleic acid sequence, which regulates expression of the nucleic acid molecule. Preferred Host Cell: The host cell is preferably a mammalian host cell.

ACTIVITY - Virucide; Hepatotropic; Cardiant; Hypotensive; Antianginal; Cytostatic; Neuroprotective; Nootropic; Antiparkinsonian; Anticonvulsant; Immunosuppressive; Antiinflammatory; Dermatological. Preferred Vector: The vector of comprises a nucleic acid sequence, which regulates expression of the nucleic acid molecule.

MECHANISM OF ACTION - Protein Kinase; Gene Therapy.

USE - The protein kinase or the nucleic acid encoding the protein is useful for modulating cellular growth, differentiation and/or development, and for modulating cellular metabolic pathways, particularly for regulating one or more proteins involved in growth and metabolism. (I) is also useful as primers or hybridization probes for detecting protein kinase-encoding nucleic acids, in tissue typing, chromosome mapping or forensic biology. These are also useful for treating viral infections (e.g. hepatitis B), cellular growth related disorders (e.g. heart failure, hypertension, atrial fibrillation, dilated and idiopathic cardiomyopathy or angina), proliferative or differentiative disorders such as cancer (e.g. liver, melanoma, prostate, cervical, breast, colon or sarcoma), disorders related with programmed cell death (e.g. Alzheimer's disease, Parkinson's disease or epilepsy), or autoimmune disorders (e.g. systemic lupus erythematosus).

ADMINISTRATION - Dosage is 0.001-30 mg/kg, preferably 1-10 mg/kg body weight. Administration can be through parenteral (e.g. intravenous, intradermal, subcutaneous), oral (e.g. inhalation), transdermal (topical), transmucosal or rectal routes.

EXAMPLE - No suitable example given.(50 pages)

=> d his

(FILE 'HOME' ENTERED AT 12:20:43 ON 07 APR 2006)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:21:52 ON 07 APR 2006

```
L1      1 S "14171 KINASE?"
L2      1434878 S KINASE?
L3      7629584 S CLON? OR EXPRESS? OR RECOMBINANT
L4      73 S "T-P" MOTIF
L5      49 S L2 AND L4
L6      12 DUP REM L5 (37 DUPLICATES REMOVED)
L7      947418 S (MODULAT? OR ACTIVAT? OR INHIBIT? ) AND L2
L8      42 S "14171"
L9      2 S L7 AND L8
```

=> e libermann rosanna

```
E1      3      LIBERMANA/BI
E2      54     LIBERMANN/BI
E3      0 --> LIBERMANN ROSANNA/BI
E4      1      LIBERMANS/BI
E5      1      LIBERMEISTER/BI
E6      1      LIBERMORE/BI
E7      1      LIBERNAN/BI
E8      5      LIBERNETICAL/BI
E9      1      LIBERNIELLA/BI
E10     282     LIBERO/BI
E11     88     LIBEROBACTER/BI
E12     1      LIBEROBACTERIA/BI
```

=> e libermann/au

E1	1	LIBERMANIS OLAFS/AU
E2	3	LIBERMANMAXE M/AU
E3	2 -->	LIBERMANN/AU
E4	1	LIBERMANN A/AU
E5	1	LIBERMANN A D/AU
E6	1	LIBERMANN A M/AU
E7	1	LIBERMANN A N/AU
E8	2	LIBERMANN B/AU
E9	1	LIBERMANN B E/AU
E10	1	LIBERMANN B M/AU
E11	1	LIBERMANN BERND/AU
E12	23	LIBERMANN C/AU

=> e kapeller r/au

E1	184	KAPELLER P/AU
E2	36	KAPELLER PETER/AU
E3	103 -->	KAPELLER R/AU
E4	4	KAPELLER REGINE/AU
E5	1	KAPELLER ROSAN/AU
E6	44	KAPELLER ROSANA/AU
E7	4	KAPELLER ROSANNA/AU
E8	2	KAPELLER RUDOLF/AU
E9	2	KAPELLER S/AU
E10	3	KAPELLER SHE A M/AU
E11	1	KAPELLER W/AU
E12	1	KAPELLERADLER REGINE/AU

=> s e3

L10 103 "KAPELLER R"/AU

=> d his

(FILE 'HOME' ENTERED AT 12:20:43 ON 07 APR 2006)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:21:52 ON 07 APR 2006

L1	1 S	"14171 KINASE?"
L2	1434878 S	KINASE?
L3	7629584 S	CLON? OR EXPRESS? OR RECOMBINANT
L4	73 S	"T-P" MOTIF
L5	49 S	L2 AND L4
L6	12	DUP REM L5 (37 DUPLICATES REMOVED)
L7	947418 S	(MODULAT? OR ACTIVAT? OR INHIBIT?) AND L2
L8	42 S	"14171"
L9	2 S	L7 AND L8
		E LIBERMANN ROSANNA
		E LIBERMANN/AU
		E KAPELLER R/AU
L10	103 S	E3

=> s l8 and l10

L11 0 L8 AND L10

=> d his

(FILE 'HOME' ENTERED AT 12:20:43 ON 07 APR 2006)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:21:52 ON 07 APR 2006

L1	1 S	"14171 KINASE?"
L2	1434878 S	KINASE?
L3	7629584 S	CLON? OR EXPRESS? OR RECOMBINANT
L4	73 S	"T-P" MOTIF

L5 49 S L2 AND L4
L6 12 DUP REM L5 (37 DUPLICATES REMOVED)
L7 947418 S (MODULAT? OR ACTIVAT? OR INHIBIT?) AND L2
L8 42 S "14171"
L9 2 S L7 AND L8
E LIBERMANN ROSANNA
E LIBERMANN/AU
E KAPPELLER R/AU
L10 103 S E3
L11 0 S L8 AND L10

	Issue Date	Page s	Document ID	Title
1	20040311	62	US 2004004830 5 A1	14171 Protein kinase, a novel human protein kinase and uses thereof
2	20031007	50	US 6630335 B1	14171 protein kinase, a novel human protein kinase and uses thereof

	Issue Date	Pages	Document ID	Title
1	20060406	95	US 2006007552 2 A1	Genes and uses for plant improvement
2	20060330	191	US 2006006838 2 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
3	20060316	157	US 2006005766 7 A1	Isolated human transporter proteins nucleic acid molecules encoding human transporter proteins and uses thereof
4	20060309	342	US 2006005185 0 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
5	20060119	83	US 2006001417 7 A1	Stable protein storage and stable nucleic acid storage in recoverable form
6	20051208	450	US 2005027167 6 A1	Inducing cellular immune responses to human immunodeficiency virus-1 using peptide and nucleic acid compositions
7	20051103	540	US 2005024483 4 A1	Single nucleotide polymorphisms in genes

8	20051006	121	US 2005022143 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
---	----------	-----	--------------------------	---

	Issue Date	Pages	Document ID	Title
9	20051006	397	US 2005022131 1 A1	Isolated human transporter proteins nucleic acid molecules encoding human transporter proteins and used thereof
10	20050901	79	US 2005019164 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
11	20050901	605	US 2005019133 1 A1	Medical implants and anti-scarring agents
12	20050825	109	US 2005018661 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
13	20050825	605	US 2005018372 8 A1	Medical implants and anti-scarring agents
14	20050818	605	US 2005018197 7 A1	Medical implants and anti-scarring agents
15	20050818	55	US 2005018136 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
16	20050818	605	US 2005018101 1 A1	Medical implants and anti-scarring agents
17	20050818	605	US 2005018100 8 A1	Medical implants and anti-scarring agents

	Issue Date	Pages	Document ID	Title
18	20050811	603	US 2005017722 5 A1	Medical implants and anti-scarring agents
19	20050811	605	US 2005017566 3 A1	Medical implants and anti-scarring agents
20	20050804	151	US 2005017041 3 A1	Isolated human ion channel proteins, nucleic acid molecules encoding human ion channel proteins, and uses thereof
21	20050728	605	US 2005016548 8 A1	Medical implants and anti-scarring agents
22	20050728	45	US 2005016521 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
23	20050728	38	US 2005016429 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
24	20050721	91	US 2005015831 2 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

25	20050714	59	US 2005015419 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
----	----------	----	--------------------------	--

	Issue Date	Page s	Document ID	Title
26	20050707	605	US 2005014915 8 A1	Medical implants and anti-scarring agents
27	20050707	605	US 2005014908 0 A1	Medical implants and anti-scarring agents
28	20050630	605	US 2005014381 7 A1	Medical implants and anti-scarring agents
29	20050623	40	US 2005013651 4 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
30	20050623	214	US 2005013647 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins and uses thereof
31	20050616	191	US 2005013088 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
32	20050616	254	US 2005013021 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
33	20050609	215	US 2005012585 2 A1	Novel kinases

	Issue Date	Page s	Document ID	Title
34	20050609	92	US 2005012398 2 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
35	20050602	36	US 2005011860 1 A1	Enhancer sequence of the 5-aminolevulinic acid synthase gene
36	20050526	70	US 2005011268 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human, transporter proteins, and uses thereof
37	20050526	248	US 2005011266 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
38	20050519	42	US 2005010667 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
39	20050428	97	US 2005008995 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Pages	Document ID	Title
40	20041209	144	US 2004024824 8 A1	Isolated human transporters proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
41	20041209	35	US 2004024811 2 A1	Isolated human transporter proteins nucleic acid molecules encoding human transporter proteins and uses thereof
42	20041209	211	US 2004024759 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
43	20041125	229	US 2004023509 3 A1	Isolated human transporter proteins nucleic acid molecules encoding human transporter proteins and uses thereof
44	20041118	64	US 2004022978 2 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
45	20041118	95	US 2004022931 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Pages	Document ID	Title
46	20041118	248	US 2004022930 4 A1	ISOLATED HUMAN GLUTAMATE RECEPTOR DNA
47	20041028	46	US 2004021485 5 A1	Cannabinoid receptor ligands and uses thereof
48	20040930	57	US 2004019289 0 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
49	20040930	103	US 2004019182 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
50	20040923	95	US 2004018552 7 A1	Isolated human transporter proteins nucleic acid molecules encoding human transporter proteins and uses thereof
51	20040826	47	US 2004016649 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
52	20040729	69	US 2004014773 2 A1	Novel human G- protein coupled receptor, HGPRBY9, expressed highly in brain and testes

	Issue Date	Pages	Document ID	Title
53	20040729	62	US 2004014688 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
54	20040701	319	US 2004012744 6 A1	Oligonucleotide mediated inhibition of hepatitis B virus and hepatitis C virus replication
55	20040624	65	US 2004012221 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
56	20040610	170	US 2004011093 8 A1	Proteins, genes and their use for diagnosis and treatment of schizophrenia
57	20040603	50	US 2004010677 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
58	20040527	256	US 2004010238 9 A1	Nucleic acid-mediated treatment of diseases or conditions related to levels of vascular endothelial growth factor receptor (VEGF-R)

	Issue Date	Page s	Document ID	Title
59	20040429	44	US 2004008203 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
60	20040422	357	US 2004007756 5 A1	Enzymatic nucleic acid-mediated treatment of ocular diseases or conditions related to levels of vascular endothelial growth factor receptor (VEGF-R)
61	20040408	154	US 2004006752 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
62	20040318	312	US 2004005338 5 A1	Crystal structure
63	20040311	322	US 2004004835 0 A1	Crystal of bacterial core RNA polymerase with rifampicin and methods of use thereof
64	20040311	62	US 2004004830 5 A1	14171 Protein kinase, a novel human protein kinase and uses thereof
65	20040304	515	US 2004004337 8 A1	Methods of identifying modulators of bromodomains

	Issue Date	Page s	Document ID	Title
66	20040205	95	US 2004002332 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
67	20040115	289	US 2004000948 8 A1	Nucleic acids, proteins, and antibodies
68	20031002	68	US 2003018638 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
69	20030925	174	US 2003018088 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
70	20030911	68	US 2003017081 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
71	20030911	92	US 2003017077 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Pages	Document ID	Title
72	20030904	43	US 2003016652 2 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
73	20030904	82	US 2003016618 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins and uses thereof
74	20030904	63	US 2003016615 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
75	20030904	63	US 2003016615 4 A1	Isolated human ion channel proteins, nucleic acid molecules encoding human ion channel proteins, and uses thereof
76	20030904	96	US 2003016615 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
77	20030828	58	US 2003016227 4 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Page s	Document ID	Title
78	20030821	80	US 2003015708 2 A1	Methods and compositions for treating cancer using 140, 1470, 1686, 2089, 2427, 3702, 5891, 6428, 7181, 7660, 25641, 69583, 49863, 8897, 1682, 17667, 9235, 3703, 14171, 10359, 1660, 1450, 18894, 2088, 32427, 2160, 9252, 9389, 1642, 85269, 10297, 1584, 9525, 14124, 4469, 8990, 2100, 9288, 64698, 10480, 20893, 33230, 1586, 9943, 16334, 68862, 9011, 14031, 6178, 21225, 1420, 32236, 2099, 2150, 26583, 2784, 8941, 9811, 27444, 50566 or 66428 molecules
79	20030807	14	US 2003014908 2 A1	INHIBITORS OF THE ANANDAMIDE TRANSPORTER AS ANALGESIC AGENTS
80	20030807	40	US 2003014845 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
81	20030807	95	US 2003014836 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Page s	Document ID	Title
82	20030731	42	US 2003014368 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
83	20030731	46	US 2003014368 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
84	20030731	67	US 2003014362 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
85	20030724	63	US 2003013882 0 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
86	20030508	66	US 2003008729 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

87	20030501	142	US 2003008273 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
----	----------	-----	--------------------------	--

	Issue Date	Page s	Document ID	Title
88	20030424	121	US 2003007777 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
89	20030424	61	US 2003007775 0 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
90	20030410	306	US 2003006861 5 A1	Polypeptides that bind HIV gp120 and related nucleic acids, antibodies, compositions, and methods of use
91	20030320	214	US 2003005449 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins and uses thereof
92	20030206	69	US 2003002774 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
93	20030130	109	US 2003002230 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Pages	Document ID	Title
94	20030123	92	US 2003001754 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
95	20030116	52	US 2003001315 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
96	20030102	45	US 2003000354 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
97	20021219	79	US 2002019276 2 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
98	20021219	72	US 2002019276 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
99	20021114	506	US 2002016863 8 A1	Compositions, kits, and methods for identification, assessment, prevention, and therapy of human prostate cancer

	Issue Date	Page s	Document ID	Title
100	20021010	34	US 2002014730 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
101	20021010	321	US 2002014714 0 A1	Nucleic acids, proteins, and antibodies
102	20021003	193	US 2002014293 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
103	20021003	242	US 2002014238 3 A1	Isolated nucleic acid molecules encoding human transport proteins
104	20021003	114	US 2002014238 1 A1	ISOLATED NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS, AND USES THEREOF
105	20021003	380	US 2002014237 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
106	20021003	53	US 2002014237 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Pages	Document ID	Title
107	20021003	173	US 2002014230 3 A1	Proteins, genes and their use for diagnosis and treatment of Schizophrenia
108	20020926	36	US 2002013712 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
109	20020919	139	US 2002013229 2 A1	NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS
110	20020912	360	US 2002012764 4 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
111	20020829	154	US 2002011951 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
112	20020829	78	US 2002011946 2 A1	Molecular toxicology modeling
113	20020822	58	US 2002011516 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Page s	Document ID	Title
114	20020822	150	US 2002011513 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
115	20020815	50	US 2002011085 2 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
116	20020808	45	US 2002010672 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
117	20020801	64	US 2002010333 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
118	20020801	207	US 2002010311 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

119	20020801	42	US 2002010263 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
-----	----------	----	--------------------------	--

	Issue Date	Pages	Document ID	Title
120	20020627	74	US 2002008219 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
121	20020627	96	US 2002008219 0 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
122	20020627	101	US 2002008167 8 A1	ISOLATED NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS, AND USES THEREOF
123	20020627	41	US 2002008165 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
124	20020627	45	US 2002008164 9 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
125	20020627	213	US 2002008164 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

	Issue Date	Page s	Document ID	Title
126	20020620	59	US 2002007675 0 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
127	20020613	56	US 2002007248 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
128	20020530	63	US 2002006482 1 A1	Isolated human ion channel proteins, nucleic acid molecules encoding human ion channel proteins, and uses thereof
129	20020425	95	US 2002004878 7 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
130	20020418	100	US 2002004516 6 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
131	20020411	201	US 2002004210 0 A1	Isolated human ion channel proteins, nucleic acid molecules encoding human ion channel proteins, and uses thereof

	Issue Date	Page s	Document ID	Title
132	20020404	67	US 2002003999 1 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
133	20020328	45	US 2002003754 8 A1	Isolated human transporter proteins, nucleic acid molecules incoding hum an transporter proteins, and uses thereof
134	20020314	69	US 2002003180 0 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
135	20020307	36	US 2002002891 5 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
136	20020307	76	US 2002002877 3 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof

137	20020214	49	US 2002001902 8 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
-----	----------	----	--------------------------	--

	Issue Date	Pages	Document ID	Title
138	20020117	75	US 20020006618 A1	Methods for using 20893, a human protein kinase
139	20011213	46	US 20010051361 A1	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
140	20060131	58	US 6991920 B2	Isolated human transporter proteins, nucleic acid molecules, encoding human transporter proteins, and uses thereof
141	20050412	207	US 6878808 B2	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins and uses thereof
142	20041214	241	US 6830900 B2	Isolated human glutamate receptor DNA
143	20040831	292	US 6783961 B1	Expressed sequence tags and encoded human proteins
144	20040831	137	US 6783930 B1	Development of novel anti-microbial agents based on bacteriophage genomics
145	20040608	872	US 6747137 B1	Nucleic acid sequences relating to Candida albicans for diagnostics and therapeutics
146	20040427	465	US 6727063 B1	Single nucleotide polymorphisms in genes
147	20031028	312	US 6639063 B1	EST's and encoded human proteins

	Issue Date	Pages	Document ID	Title
148	20031007	50	US 6630335 B1	14171 protein kinase, a novel human protein kinase and uses thereof
149	20030513	93	US 6562593 B2	Isolated human transporter proteins, nucleic acid molecules encoding human transporter proteins, and uses thereof
150	20021224	238	US 6498022 B2	Isolated nucleic acid molecules encoding human carbonate transporter proteins, and uses thereof
151	19980203	132	US 5714503 A	Compounds and methods for inhibition of HIV and related viruses
152	19970819	130	US 5658907 A	Compounds and methods for inhibition of HIV and related viruses
153	19970114	130	US 5593993 A	Method for inhibition of HIV related viruses
154	19941011	13	US 5354744 A	Side chain unsaturated 1 alpha-hydroxyvitamin D analogs
155	19931005	14	US 5250523 A	Side chain unsaturated 1.alpha.-hydroxyvitamin D homologs
156	19930511	16	US 5210029 A	Method of producing interleukin-2
157	19900619	8	US 4935356 A	Production of interleukin-2
158	19870721	29	US 4681848 A	Novel peptide and use thereof

	Issue Date	Page s	Document ID	Title
1	20040311	62	US 2004004830 5 A1	14171 Protein kinase, a novel human protein kinase and uses thereof
2	20020117	75	US 2002000661 8 A1	Methods for using 20893, a human protein kinase
3	20031007	50	US 6630335 B1	14171 protein kinase, a novel human protein kinase and uses thereof

	L #	Hits	Search Text
1	L1	2	"14171" adj kinase\$2
2	L2	1510 328	modulat\$2 or inhibit\$3 or activat\$3
3	L3	8297 66	clon\$3 or express\$3 or recombinant
4	L4	1	"t-P" adj moti\$3
5	L5	0	"t-P moti\$3"
6	L6	7001 3	kinase\$2
7	L7	1342 70	12 same 13
8	L8	1342 70	13 same 17
9	L9	372	"14171"
10	L10	160	18 and 19
11	L11	158	human and 110
12	L12	135	KAPELLER-LIBERMANN KAPELLER-ROSANA
13	L13	3	19 and 112